

Isometric stability property of certain Banach spaces

ALEXANDER KOLDOBSKY

Abstract. Let E be one of the spaces $C(K)$ and L_1 , F be an arbitrary Banach space, $p > 1$, and (X, σ) be a space with a finite measure. We prove that E is isometric to a subspace of the Lebesgue-Bochner space $L_p(X; F)$ only if E is isometric to a subspace of F . Moreover, every isometry T from E into $L_p(X; F)$ has the form $Te(x) = h(x)U(x)e$, $e \in E$, where $h : X \rightarrow R$ is a measurable function and, for every $x \in X$, $U(x)$ is an isometry from E to F .

Mailing address:.

Till August 1, 1993:
 Department of Mathematics
 University of Missouri-Columbia
 Columbia, MO 65211, USA

After August 1, 1993:
 Division of Mathematics, Computer Science and Statistics
 University of Texas at San Antonio
 San Antonio, TX 78249, USA

AMS classification:. 46B04, 47B80

Key words:. isometries, Lebesgue-Bochner spaces, random operators

Running title:. Isometric embeddings

Isometric stability property of certain Banach spaces

ALEXANDER KOLDOBSKY¹

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MISSOURI-COLUMBIA
COLUMBIA, MO 65211

1.Introduction.

Let E and F be Banach spaces, $p \geq 1$, (X, σ) be a finite measure space, and $L_p(X, F)$ be the Lebesgue-Bochner space of (equivalence classes of) strongly measurable functions $f : X \mapsto F$ with

$$\|f\|^p = \int_X \|f(x)\|^p d\sigma(x) < \infty.$$

We show that if $E = C(K)$ (with K being a compact metric space) or $E = L_1$ then the space E can be isometric to a subspace of $L_p(X, F)$ with $p > 1$ only if E is isometric to a subspace of F .

The isomorphic version of this result has been proved for the spaces $E = c_0$ (S.Kwapień [5] and J.Bourgain [1]), $E = l_1$ (G.Pisier [7]) and $E = l_\infty$ (J.Mendoza [6], see also [2] for generalizations to Kothe spaces of vector valued functions). E.Saab and P.Saab [9] have proved the isomorphic version for the space $E = L_1$ under the additional assumption that F is a dual space.

For any $1 \leq p \leq q \leq r$, the space L_q is isometric to a subspace of $L_p(X, L_r)$ (see [8]). On the other hand, if $r > 2, r \neq q, q \neq 2$ the space L_q is not isomorphic to a subspace of L_r . Thus, both isometric and isomorphic versions fail to be true in the case where $E = L_q, q > 1$. (The space L_2 is isometric to a subspace of L_p for any $p > 0$, so it is isometric to a subspace of $L_p(X, F)$ for any space F)

Besides proving the above mentioned result for the spaces $C(K)$ and L_1 , we completely characterize isometric embeddings of these spaces into L_p -spaces of vector valued functions.

Denote by $I(E, F)$ the set of isometries from E to F . A mapping $U : X \mapsto I(E, F)$ is called strongly measurable if, for each $e \in E$, the function $\|U(x)e\|$ is measurable on X .

¹After August 1, 1993 the author's address will be: Division of Mathematics, Computer Science and Statistics, University of Texas at San Antonio, San Antonio, TX 78249.

If the set $I(E, F)$ is non-empty then, obviously, for every strongly measurable mapping $U : X \mapsto I(E, F)$ and every function $h : X \mapsto \mathbb{R}$ with $\|h\|_{L_p(X)} = 1$, the operator $T : E \mapsto L_p(X, F)$ defined by

$$(1) \quad Te(x) = h(x)U(x)e, e \in E$$

is an isometry.

We prove below that, for $E = C(K)$ or $E = L_1$ and for an arbitrary space F , every isometry from E to $L_p(X, F)$ has the form (1). The question if all isometries from E to $L_p(X, F)$ have the form (1) makes sense and has some applications even if $E = F$. For example, if $E = F = L_q, q > 1, p > 1$, then every isometry from L_q to $L_p(X, L_q)$ has the form (1) if and only if $p \neq q, q \neq 2, q \notin (p, 2)$. This result was proved and applied to the description of isometries of Lebesgue-Bochner spaces in the paper [3].

As a consequence of the characterization of isometries from $E = C(K)$ or $E = L_1$ to $L_p(X, F)$ we obtain the following result on random operators. Suppose that (X, σ) is a probability space, let $p > 1$, $L(E, F)$ be the space of linear operators from E to F and consider a random operator $S : X \mapsto L(E, F)$ which is an isometry in average, i.e. S is strongly measurable and, for every $e \in E$,

$$\|e\|^p = \int_X \|S(x)e\|^p d\sigma(x).$$

Then the random operator S is an isometry (up to a constant) with probability 1 (namely, operators $S(x)$ are isometries multiplied by constants for almost all $x \in X$). In fact, if we define an operator $T : E \mapsto L_p(X, F)$ by $Te(x) = S(x)e$ then T is an isometry and (1) implies the desired result. In the case $E = C(K)$, the result about random operators has been proved before [4] under the assumption that the operators $S(x)$ are bounded.

2. Main results.

The proofs are based on the following simple fact.

LEMMA 1. *Let E, F be Banach spaces, $p > 1$, (X, σ) be a finite measure space and T be an isometry from E to $L_p(X, F)$. If e, f are elements from E with $\|e\| = \|f\| = 1$ and having a common tangent functional (i.e. there exists $x^* \in E^*$ with $\|x^*\| = 1, x^*(e) = x^*(f) = 1$) then, for almost all (with respect to σ) $x \in X$, we have*

- (i) $\|Te(x)\| = \|Tf(x)\|$,
- (ii) for every $\alpha > 0$, $\|Te(x) + \alpha Tf(x)\| = \|Te(x)\| + \alpha\|Tf(x)\|$.

PROOF: Obviously, $\|e + \alpha f\| = 1 + \alpha$ for every $\alpha > 0$, and we have

$$(1 + \alpha)^p = \|e + \alpha f\|^p = \int_X \|Te(x) + \alpha Tf(x)\|^p d\sigma(x) \leq$$

$$(2) \quad \int_X (\|Te(x)\| + \alpha\|Tf(x)\|)^p d\sigma(x)$$

For $\alpha = 0$, (2) turns into an equality. Therefore, we get a correct inequality if we take in both sides of (2) the right-hand derivatives at the point $\alpha = 0$ and apply Holder's inequality:

$$1 \leq \int_X \|Te(x)\|^{p-1} \|Tf(x)\| d\sigma(x) \leq \left(\int_X \|Te(x)\|^p d\sigma(x) \right)^{\frac{p-1}{p}} \left(\int_X \|Tf(x)\|^p d\sigma(x) \right)^{1/p} = 1$$

From the conditions for equality in Holder's inequality, we conclude that, for almost all (with respect to σ) $x \in X$, $\|Te(x)\| = c\|Tf(x)\|$ where c is a constant. Further,

$$1 = \int_X \|Te(x)\|^p d\sigma(x) = \int_X c^p \|Tf(x)\|^p d\sigma(x) = c^p$$

and, hence, $c = 1$.

It is clear now that (2) is, in fact, an equality. Hence, for almost all $x \in X$, we have

$$(3) \quad \|Te(x) + \alpha Tf(x)\| = \|Te(x)\| + \alpha\|Tf(x)\|$$

for every $\alpha > 0$ and the proof is complete. ■

Now we are able to prove the main result.

THEOREM 1. *Let $p > 1$, K be a compact metric space, $(X, \sigma), (Y, \nu)$ be spaces with finite measures, and F be an arbitrary Banach space. Let E be either the space $L_1 = L_1(Y, \nu)$ or any subspace of $C(K)$ containing the function $1(k) \equiv 1$. Then*

(i) *If E is isometric to a subspace of $L_p(X; F)$ then E is isometric to a subspace of F and the set $I(E, F)$ is non-empty.*

(ii) *If T is an isometry from E into $L_p(X; F)$ then there exist a measurable function $h : X \rightarrow R$ and a strongly measurable mapping $U : X \rightarrow I(E, F)$ such that $Te(x) = h(x)U(x)e$ for every $e \in E$.*

PROOF: We start with the case $E = L_1$.

Any two functions e and f from L_1 with disjoint supports in Y have a common tangent functional so we can apply Lemma 1 to any pair of normalized functions with disjoint supports.

Decompose the set Y into two parts $Y = Y_1 \cup Y_2, Y_1 \cap Y_2 = \emptyset, \nu(Y_i) > 0, i = 1, 2$. Fix a function $e_0 \in L_1(Y_1), \|e_0\| = 1$ and put $h(x) = \|Te_0(x)\|, x \in X$.

Let $f_k, k \in N$ be a sequence of linearly independent functions with supports in Y_2 such that their linear span is dense in $L_1(Y_2)$. Denote by D the set of linear combinations of functions f_k with rational coefficients. Given fixed representatives Tf_k from the corresponding equivalence classes of functions from the space $L_p(X; F)$, define an operator $T(x) : D \mapsto F$ for every $x \in X$ by $T(x)(\sum \lambda_i f_i) = \sum \lambda_i Tf_i(x), \lambda_i \in Q$.

It follows from the statement (i) of Lemma 1 and the fact that D is countable that there exists a set $X_0 \subset X$ with $\sigma(X \setminus X_0) = 0$ such that, for every $x \in X_0$ and every $f \in D$,

$$\|T(x)f\| = \|Tf(x)\| = \|Te_0(x)\| \|f\| = h(x)\|f\|$$

Hence, for every $x \in X_0$, either $h(x) = 0$ or the operator $U_2(x) = T(x)/h(x)$ is an isometry from D to F .

The operators $U_2(x)$ can be uniquely extended to isometries on the whole space $L_1(Y_2)$. In fact, given $a \in L_1(Y_2)$ and a sequence $a_k \rightarrow a, a_k \in D$, put $U_2(x)(a) = \lim_{k \rightarrow \infty} U_2(x)(a_k)$.

Further, for any $a \in L_1(Y_2)$,

$$\begin{aligned} \|a_k - a\|^p &= \int_X \|Ta_k(x) - Ta(x)\|^p d\sigma(x) = \\ (4) \quad &\int_{h(x)=0} \|Ta(x)\|^p d\sigma(x) + \int_{h(x) \neq 0} \|h(x)U_2(x)a_k - Ta(x)\|^p d\sigma(x) \rightarrow 0 \end{aligned}$$

as $k \rightarrow \infty$. Therefore, $Ta(x) = 0$ for almost all $x \in X$ with $h(x) = 0$, and $Ta(x) = h(x)U_2(x)a$ for almost all $x \in X$ (if $h(x) = 0$ we put $U_2(x) = 0$.)

Similarly, for almost all $x \in X$, we find isometries $U_1(x)$ from $L_1(Y_1)$ to F such that $Tb(x) = h(x)U_1(x)b$ for every $b \in L_1(Y_2)$.

Consider an arbitrary function $f \in L_1(Y)$. This function can be uniquely represented as a sum $f = f_1 + f_2$ of functions $f_1 \in L_1(Y_1)$ and $f_2 \in L_1(Y_2)$. For all $x \in X$ with $h(x) \neq 0$, define operators $U(x)$ from $L_1(Y)$ to F by $U(x)f = U_1(x)f_1 + U_2(x)f_2$. By the statement (ii) of Lemma 1, for almost every x with $h(x) \neq 0$,

$$\|U(x)f\| = (1/h(x))\|Tf_1(x) + Tf_2(x)\| = \|U_1(x)f_1\| + \|U_2(x)f_2\| = \|f_1\| + \|f_2\| = \|f\|$$

Thus, operators $U(x)$ are isometries for almost all $x \in X$ with $h(x) \neq 0$. In particular, the set $I(L_1, F)$ is non-empty. Fix any $U \in I(L_1, F)$ and put $U(x) = U$ for every x with $h(x) = 0$.

To prove the second statement of Theorem 1 note that, for every $f \in L_1(Y)$, we have $Tf(x) = Tf_1(x) + Tf_2(x) = h(x)(U_1(x)f_1 + U_2(x)f_2) = h(x)U(x)f$ for almost all $x \in X$, so Tf and $h(x)U(x)f$ are equal elements of the space $L_p(X, F)$.

Now let E be a subspace of $C(K)$ containing the function $1(k) \equiv 1$.

Any function $e \in C(K)$ has a common tangent functional either with the function 1 or with the function -1 . Setting, correspondingly, $f = 1$ or $f = -1$ in Lemma 1 we obtain that, for an arbitrary $e \in E$, $\|Te(x)\| = \|T1(x)\|\|e\|$ for almost all $x \in X$. Let $h(x) = \|T1(x)\|$.

Let $e_k, k \in N$ be a sequence of linearly independent functions from E such that their linear span is dense in E and denote by D the set of linear combinations of functions e_k with rational coefficients. Given fixed representatives Te_k from the corresponding equivalence classes of functions from the space $L_p(X; F)$, define an operator $T(x)$ on D for every $x \in X$ by $T(x)(\sum \lambda_i e_i) = \sum \lambda_i Te_i(x), \lambda_i \in R$.

Since the set D is countable there exists a set $X_0 \subset X$ with $\sigma(X \setminus X_0) = 0$ such that, for every $x \in X_0$ and every $e \in D$, $\|T(x)e\| = \|Te(x)\| = h(x)\|e\|$. Hence, for every $x \in X_0$, either $h(x) = 0$ or the operator $U(x) = T(x)/h(x)$ is an isometry from D to F .

The operators $U(x)$ can be uniquely extended to isometries on the whole space E , therefore, we have proved the first statement of Theorem 1.

Now an argument similar to (4) proves the second statement. ■

Remark. If $p = 1$ the statement of Theorem 1 is not true. For instance, the two-dimensional space l_∞^2 is isometric to a subspace of $L_1([0, 1])$. Thus, for any Banach space F , l_∞^2 is isometric to a subspace of $L_1([0, 1]; F)$, and Theorem 1 would have implied that l_∞^2 is isometric to a subspace of any Banach space.

References.

1. Bourgain, J.: An averaging result for c_0 -sequences, Bull. Soc. Math. Belgique, Vol.B 30(1978), 83-87
2. Emmanuele, G.: Copies of l_∞ in Kothe spaces of vector valued functions, Illinois J. Math. 36(1992), 293-296
3. Koldobsky, A.: Isometries of $L_p(X; L_q)$ and equimeasurability, Indiana Univ. Math. J. 40(1991), 677-705
4. Koldobsky, A.: Measures on spaces of operators and isometries, J. Soviet Math. 42(1988), 1628-1636
5. Kwapien, S.: On Banach spaces containing c_0 , Studia Math. 52(1974), 187-188
6. Mendoza, J.: Copies of l_∞ in $L^p(\mu; X)$, Proc. Amer. Math. Soc. 109(1990), 125-127
7. Pisier, G.: Une propriete de stabilite de la classe des espaces ne contenant pas l_1 , C.R. Acad. Sci. Paris Ser A 86(1978), 747-749
8. Raynaud, Y.: Sous espaces l^r et geometrie des espaces $L^p(L^q)$ et L^ϕ , C.R. Acad. Sci. Paris, Ser.I 301(1985), 299-302
9. Saab, E. and Saab, P.: On stability problems of some properties in Banach spaces, Function spaces (K.Jarosz, editor), Lect. Notes in Pure and Appl. Math. 136(1992), 367-394